



**ISSX**

International Society for the  
Study of Xenobiotics



# 9th International ISSX Meeting

**September 4 — 8, 2010**  
**Istanbul, Turkey**

## PROGRAM & ABSTRACTS

ISSX would like to thank:

Turkish Ministry of Culture and Tourism

Co-Sponsor of the Short  
Course Programs:



EUROPEAN FEDERATION  
FOR PHARMACEUTICAL  
SCIENCES



Middle East Technical University  
Orta Doğu Teknik Üniversitesi



Türkiye Bilimler Akademisi  
Turkish Academy of Sciences





### P152. SEIZUROGENIC EFFECTS OF LOW-DOSE NALOXONE IN TRAMADOL OVERDOSE

Esmail Farzaneh<sup>1</sup>, Babak Mostafazadeh<sup>2</sup>, Farnaz nasl Seraji<sup>1</sup> and Iraj sayad Rezai<sup>1</sup>

<sup>1</sup>Medical Toxicology and Forensic Medicine, Ardebil University of Medical Sciences, Ardabil, Iran, Ardabil, Iran,

<sup>2</sup>Medical Toxicology & Forensic Medicine, Shaheed Beheshty University Of Medical Sciences, Tehran, Iran

**Objective:** Tramadol is used in treatment of moderate to severe pain. Nowadays tramadol overdose is one of the common emergencies. Naloxone is an antagonist which is used as a first step of treatment in these patients. This study was designed to evaluate the seizurogenic effects of naloxone in tramadol overdose. **Methods:** 124 patients with the diagnosis of tramadol overdose were divided to receive low-doses of intravenous naloxone (0.8 mg, case group) or just supportive cares (control group). All patients in case and control groups were observed by a single emergency resident and followed for 1.5 hours to document the happening of seizures. **Results:** In the naloxone group, incidence of seizure was higher than in control group. The possibility of seizure occurrence was significantly higher in naloxone group than the control group ( $P < 0.05$ ). **Conclusion:** Naloxone induced a seizurogenic effect in patients with tramadol overdose. This finding could be considered in the management of patients with tramadol overdose.

#### References

1. Rehni AK, Singh I, Kumar M. Tramadol-induced seizurogenic effect: a possible role of opioid-dependent aminobutyric acid pathway. *Basic Clin Pharmacol Toxicol* 2008; 103: 262-66.
2. Ashish K. Rehni, Thakur Gurjeet Singh, Nirmal Singh<sup>2</sup> and Sandeep Arora<sup>1</sup> Tramadol-induced seizurogenic effect: a possible role of opioid-dependent histamine ( $H_1$ ) receptor activation-linked mechanism. *Naunyn Schmiedeberg's Arch Pharmacol* 2009 DOI: 10.1007/s00210-009-0476-y
3. Yang L, Li F, Ge W, Mi C, Wang R, Ruopeng Sun R. Protective effects of naloxone in two-hit seizure model. *Epilepsia* 2009 (In Press).

Table 1. Demographic characteristics of 124 patients with tramadol overdose

Variable	Case (N = 62)	Control (N = 62)	p-value
Age(year)	26.33	29.46	0.5
Male	59	55	0.18
Female	3	7	—
Single	35	27	0.15
Married	27	35	

P-value less than 0.05 considered significant

Table 2: Comparison of past medical history of the patients with tramadol overdose between the two studied groups (N = 124).

	Case (N = 62)	Control (N = 62)	p-value
Former drug overdose	24	34	0.07
Former tramadol overdose	8	7	0.78
History of seizure	3	5	0.71
CNS* disease	3	5	0.38
Alcoholism	0	1	1.00
Cigarette smoking	52	45	0.12

\* Central Nervous System

### P153. SEQUENTIAL ARABINOSYLCYTOSIN WITH OR WITHOUT FLUDARABINE IN PARACMASTIC PATIENTS OF ACUTE MYELOID LEUKEMIA

Ai-Bin Liang<sup>1</sup>, Zhang Hong<sup>1</sup>, Yu Fang, Ying Li, De-Qiu Zhu<sup>1</sup>, Fang-Yuan Chen<sup>2</sup>, Hui-Na Lu<sup>1</sup>, Jian-Fei Fu<sup>1</sup>, Lan-Jun Bo<sup>1</sup>

<sup>1</sup>Clinical Pharmacology, Tongji Hospital of Tongji University, Shanghai, China, <sup>2</sup>Renji Hospital, Shanghai Jiaotong University, Shanghai, China

**Objective:** The synergistic action between Fludarabine(Fa) and Arabinosylcytosin(Ara-C) has been well reported. However, this phenomena doesn't always happen in clinical, and little has been studied on how to decide whether Ara-C can be combined with Fa or not in clinical. The purpose of this study is to assess what the difference of Ara-C's behavior is with or without Fa. **Methods:** The plasma, cerebrospinal and urinal samples from two self-control groups (Group A with Ara-C only, Group B with Fa + Ara-C) were withdrawn at specific time points, and analyzed with HPLC. **Results:** The  $AUC_{0-4}$  of Group B and Group A was  $12.245 \pm 3.863$  and  $5.131 \pm 0.936$  respectively, which there existed statistics difference ( $P=0.016$ ). The Ara-C dose in Group B was  $2g/m^2 \cdot A_2$ , but the  $AUC_{0-4}$  was over double than that of Group A ( $3g/m^2 \cdot A_2$ ); It suggests that there might exist difference in the ratio of  $C_{Ara-U}$  and  $C_{Ara-C}$  and  $T_{max}$  of the two concentrations' ratio between Group A and B; Fa might conduce to the increasing of the concentration of  $C_{Ara-C}$  in cerebrospinal fluid; The ratio of  $C_{6\beta\text{-hydroxycortisol}}/C_{cortisol}$  (a noninvasive marker for CYP3A enzymes activity) was below 1.5 in the two groups, which was obviously lower than that of healthy subjects. **Conclusions:** It is indicated that there existed synergistic action between Fa and Ara-C, in which Fa conduces to increasing the concentration and AUC of Ara-C in plasma, and even increasing the Ara-C concentration in cerebrospinal fluid. The obvious change in  $T_{max}$  of